

**ERRATUM: “The QSO evolution derived from the *HBQS*
and other complete QSO surveys ¹” [ASTRON. J., 113, 1517 (1997)]**

Fabio La Franca

Dipartimento di Fisica, Università degli studi “Roma Tre”

Via della Vasca Navale 84, I-00146 Roma, Italy

Electronic mail: lafranca@amaldi.fis.uniroma3.it

and

Stefano Cristiani

Dipartimento di Astronomia, Università degli studi di Padova

Vicolo dell’Osservatorio 5, I-35122 Padova, Italy

Electronic mail: cristiani@astrpd.pd.astro.it

The equation used for the simulations of the LDLE models is incorrectly reported. The correct equation for the LDLE evolution of the break magnitude is $M_B(z) = M_B^*(2) - 2.5k \log[(1+z)/3]$ where $M_B^*(2)$ is the break magnitude at $z = 2$ and

$$\begin{aligned} \text{for } M_B \leq M_B(z) &: k = k_1 + k_2(M_B - M_B(z))e^{-z/.40} \\ \text{for } M_B > M_B(z) &: k = k_1. \end{aligned}$$

In Table 2 the values of M_B^* for the LDLE models C and M should be substituted by the values for $M_B^*(2)$. They should read, for model C, -26.3; for model M, -26.9. This change does not concern all the other models reported in Table 2 which use the standard PLE parameterization reported in the paper. We thank L. Wisotzki for turning our attention toward this point.

¹Based on data of the ESO Key-Programme “A Homogeneous Bright QSO Survey”